

RESEARCH ARTICLE

Association of *MAOA* genetic variants and resilience with psychosocial stress: A longitudinal study of Syrian refugees

Christopher J. Clukay^{1,2}, Rana Dajani³, Kristin Hadfield⁴, Jacklyn Quinlan^{1,2}, Catherine Panter-Brick⁵, Connie J. Mulligan^{1,2*}

1 Department of Anthropology, University of Florida, Gainesville, FL, United States of America, **2** Genetics Institute, University of Florida, Gainesville, FL, United States of America, **3** Department of Biology and Biotechnology, Hashemite University, Zarqa, Jordan, **4** School of Biological and Chemical Sciences, Queen Mary, University of London, London, England, **5** Department of Anthropology, Yale University, New Haven, CT, United States of America

* cmulligan@ufl.edu



OPEN ACCESS

Citation: Clukay CJ, Dajani R, Hadfield K, Quinlan J, Panter-Brick C, Mulligan CJ (2019) Association of *MAOA* genetic variants and resilience with psychosocial stress: A longitudinal study of Syrian refugees. PLoS ONE 14(7): e0219385. <https://doi.org/10.1371/journal.pone.0219385>

Editor: Soraya Seedat, Stellenbosch University, SOUTH AFRICA

Received: February 27, 2019

Accepted: June 22, 2019

Published: July 17, 2019

Copyright: © 2019 Clukay et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are reported in the manuscript and its Supporting Information files. All data are also available on Mendeley (<https://data.mendeley.com/datasets/>) doi:10.17632/xhs8tn7nkz.1.

Funding: Laboratory research was funded by the University of Florida College of Liberal Arts and Sciences (CJM). Field research was funded by Elrha's Research for Health in Humanitarian Crises (R2HC) Programme (<http://www.elrha.org/r2hc/home>), which aims to improve health outcomes by

Abstract

Early childhood trauma can have profound and lifelong effects on adult mental health and psychosocial wellbeing. Nevertheless, responses to trauma are highly variable. Genetic variants may help explain variation in responses to trauma by identifying alleles that associate with changes in mental health measures. Protective factors, such as resilience, likely also play an important role in responses to trauma. The effects of genetic variants, in combination with protective factors, on psychosocial health are not well understood, particularly in non-Western contexts. In this study, we test the relative influence of genetic variants of monoamine oxidase A (*MAOA*, a gene proposed to influence the impact of childhood trauma on adult violence and antisocial behavior), levels of resilience, and exposure to traumatic events on psychosocial stress and mental health trajectories over time. We use data from a cohort of 12-18-year-old Syrian refugees who were forcibly displaced to neighboring Jordan ($n = 399$). DNA samples and survey data on trauma exposure, resilience (CYRM-12), and psychosocial stress were collected at three time points: baseline, ~13 weeks, and ~48 weeks. Using multilevel models, we identified an association of *MAOA* variant, in males only, with symptom scores of psychosocial stress on the Perceived Stress Scale (PSS) over time ($p = 8.1 \times 10^{-4}$). We also found that resilience is strongly associated with PSS ($p = 7.9 \times 10^{-9}$), underscoring the importance of protective factors in influencing levels of psychosocial stress. Furthermore, there was an additive effect wherein the sharpest reductions in perceived psychosocial stress are seen in low-activity *MAOA* males with low trauma exposure or high resilience levels. Our results highlight the value of studies that integrate genetic and psychosocial factors to better understand complex phenotypes, such as responses to trauma in contexts of high trauma exposure.

strengthening the evidence base for public health interventions in humanitarian crises (CPB). The R2HC programme is funded equally by the Wellcome Trust and the UK Government. Additional funding was received from NSF grant DGE-1315138. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Childhood traumatic experiences are associated with poor health outcomes that can persist throughout life. In particular, children and adolescents who experience physical and emotional abuse are at increased risk for psychiatric diagnoses, especially mood, anxiety, and substance abuse disorders, as well as suicidality, violent and criminal behavior, poor physical health and increased risk for infectious disease later in life [1–7]. However, there is great variation in how children experience and recover from early trauma [8]. There is a paucity of longitudinal research conducted outside Western contexts on this issue, especially with war-affected children and adolescents. Research on refugee populations is of great importance because refugees experience a high burden of trauma, loss, and stress [9]. Understanding how war-related adversity and trauma impacts health and development is particularly critical in children and adolescents since they account for 52% of refugees worldwide [10]. Many refugees are exposed to substantial trauma during war, as well as ongoing stressors and social marginalization during and after resettlement. Nevertheless, many refugees show notable resilience in psychosocial, developmental, and health trajectories [11, 12]. Now in its seventh year, the Syrian crisis has led more than 5.68 million people to leave Syria, while a further 6.5 million Syrians are internally displaced, two-fifths of whom are under 18 years of age [13, 14]. This is the largest refugee crisis since the Second World War, the consequences of which have led scholars and humanitarian agencies to call for renewed attention to address health and developmental issues for a potentially ‘lost generation’ coping with exposures to ‘toxic stress.’ [9].

The factors that determine how children experience and recover from trauma are not well understood, but likely include both genetic and psychosocial factors [8]. The monoamine oxidase A gene (*MAOA*) has been widely studied for its role in influencing the impact of childhood trauma on adult aggressive and antisocial behavior. *MAOA* is located on the X chromosome and the encoded *MAOA* enzyme catalyzes the degradation of brain neurotransmitters serotonin, dopamine and norepinephrine, thus playing a key role in regulating behavior. A functional repeat polymorphism located approximately 1.2 kb upstream of the *MAOA* coding sequence was first identified by Sabol et al. [15], who used luciferase reporter gene fusions and transfection experiments to show that alleles with different numbers of repeat sequences were transcribed with different efficiencies in human neuroblastoma and human placental choriocarcinoma cell lines. *MAOA* deficiency was first studied in humans in five affected males who exhibited borderline mental deficiency as well as impulsive aggression, criminal violence, and hypersexual behavior [16]. In the early 2000’s, Caspi and Moffitt’s [17] landmark Dunedin Longitudinal Cohort study investigated *MAOA* genetic variants to better understand why some abused children developed antisocial behavior as adults and others did not. They determined that childhood trauma was more likely to lead to antisocial and violent behavior in males who carried a *MAOA* variant with low transcriptional activity (*MAOA-L*) [17].

Over the past decade, additional studies have identified associations between *MAOA* variants and a wide range of aggressive and antisocial behaviors including adolescent antisocial behavior [18], reduced social cooperation [19], physical aggression [20], criminal violence [21], recidivism in male violent offenders [22], and brain morphology in violent offenders with psychopathic traits [23]. Not all associations have been confirmed, however, and the first genome-wide association study on adult antisocial behavior did not find significant associations with any genetic polymorphisms, leading the authors to speculate that the relevant genes may have effect sizes which are so small that they are undetectable with their sample size [24]. In general, *MAOA-L* is the allele most often associated with increased aggressive or antisocial behaviors and the strongest associations are typically found in males [25, 26].

Interestingly, some studies have found opposite effects in females, i.e. association of the high-activity *MAOA* (*MAOA-H*) variant with increased risk of antisocial behavior and aggression [25, 27]. A study of neural activity found that activity in the amygdala and hippocampus increased with childhood stress in *MAOA-L* males and decreased in *MAOA-H* males, with the opposite pattern seen in females [27]. Another brain imaging study found increased connectivity between the amygdala and ventromedial prefrontal cortex that indicated dysregulated amygdala activation in *MAOA-L* males; this effect was not seen in females [28]. Variable levels of testosterone and the specific interaction of *MAOA-L* with high levels of testosterone [29] likely explain some of the differing effects of *MAOA* between males and females, and possibly within males as well.

The timing of trauma exposure is critical since brain structures are still developing during childhood and adolescence; disruption of different brain structures will likely impact different psychosocial outcomes [30, 31]. The critical developmental periods, and sensitive windows, for three key regions of the brain (hippocampus, amygdala, prefrontal cortex) all overlap in early and middle childhood [32, 33]. Children across a range of ages have been studied in order to investigate critical windows of trauma exposure. For instance, family adversity experienced in the first three years of life has been found to associate with increased hyperactivity at 4–7 years of age in *MAOA-L* boys and girls from the U.K. Avon Longitudinal Study of Parents and Children, UK [34]. Fewer studies have focused on adolescents, but Lee [18] found that the effect of deviant peer affiliation on antisocial behavior in participants from the National Longitudinal Study of Adolescent Health study was significantly stronger in *MAOA-H* carriers than *MAOA-L* carriers. In sum, different studies identify both *MAOA-L* and *MAOA-H* variants as being risk alleles, which may reflect ongoing brain development and exposure to trauma at different ages.

In addition to the effect of *MAOA* on aggressive and antisocial behaviors, other behaviors have been tested for association with *MAOA* in order to provide a more comprehensive perspective on *MAOA* functioning and to better understand the behaviors impacted by *MAOA*. For instance, *MAOA-L* variants have been found at slightly higher frequencies in alcoholics (both male and female) with antisocial personality disorder compared to alcoholics with anxiety and depression, suggesting that *MAOA* variation may contribute to over- vs under-reactive behaviors in the development of alcoholism [26]. *MAOA* likely also plays a role in the risk of developing stress-related behaviors like obsessive-compulsive disorder [35] as well as addictive behaviors like cigarette smoking [36]. In a study on assessment of risk in a laboratory setting, *MAOA-H* individuals showed a preference for longshot risks and were less likely to purchase insurance, suggesting that *MAOA* variation may play a role in real-life decision-making and risk-assessment [37]. Negative associations can also be informative, such as the study that found *MAOA* was not one of the nine candidate genes associated with response time on an attention task, suggesting that *MAOA* does not play a role in attentional deficits [38]. Overall, our understanding of the effect of *MAOA* on complex behaviors is incomplete, but the impact of *MAOA* is likely to be relatively small, and possibly difficult to detect [39].

Virtually no studies have investigated the influence of *MAOA* and protective factors on psychosocial and mental health outcomes. We found a single published study of *MAOA* and a protective factor: high perceived parental care was found to mitigate the effect of childhood stress in females (males were not studied) [40]. Specifically, a three-way interaction between perceived parental care, early life family stressors, and *MAOA* variant predicted impulsivity/aggression scores. The lack of attention to protective factors is puzzling given the strong effect of protective factors on mental and physical health [41–46] and the opportunity for intervention [47, 48] with potential for intergenerational effects [49, 50]. Furthermore, from an evolutionary perspective, it is unlikely to find common genetic variants whose only effect is to

increase the risk of maladaptive behaviors under adverse conditions. Pluess and Belsky have proposed that genetic variants like those at *MAOA* do not function solely as risk factors, but rather as environmental sensitivity variants whose carriers are more sensitive, or responsive, to both positive and negative exposures [51–53]. In their brain imaging study, Buckholtz et al. [54] proposed that *MAOA-L* variants may create a less stable developmental framework, e.g. amygdala dysregulation and a compensatory response of increased coupling between the amygdala and the ventromedial prefrontal cortex, that renders *MAOA-L* males more sensitive to environmental factors relative to *MAOA-H* males.

In this study, we tested for association of *MAOA* variants with different measures of psychosocial stress and mental health in a population of Syrian refugee youth who have experienced high levels of war-related trauma. We also investigated the protective aspects of a contextually-specific measure of resilience that measures individual and collective strengths, resources, and capabilities [55]. The goal in studying a range of measures was to identify outcomes other than aggressive and antisocial behavior that associate with *MAOA*. We predicted there would be an association of *MAOA* genetic variants, either directly or through an interaction with trauma exposure or resilience, with psychosocial outcomes. Specifically, we hypothesized that *MAOA-L* individuals would show increased change in the measured outcomes relative to *MAOA-H* carriers and that this effect would be limited to males, based on findings of dysregulated amygdala activation and proposed increased environmental sensitivity in *MAOA-L* males [28, 56].

Materials and methods

Study design

Our study population was a cohort of 399 12–18-year-old Syrian refugees in Jordan. Over 671,000 Syrians have taken refuge in Jordan [14]. Participating adolescents were sampled from four sites in northern Jordan (Irbid, Jarash, Mafraq, Zarqa), urban centers which have been heavily affected by the Syrian crisis. Youth in our study were participating in an eight-week stress attunement intervention (*Advancing Adolescents*; Arabic: *Nubader*), developed by Mercy Corps for war-affected adolescents in Jordan, Lebanon, Iraq, Syria and Turkey. Buccal samples and data on trauma exposure, resilience, stress, and mental health were collected at three time points: pre-intervention baseline (T1), after ~13 weeks (T2), and after ~48 weeks (T3). Refugees at these four study sites were from disadvantaged families and most were displaced from the cities of Damascus, Daraa, Homs, and surrounding towns in Syria. In order to control for possible population structure created by study sites having different proportions of refugees from different areas of Syria, study site was included as a covariate in all analyses. Participants were sampled in two waves: in the first wave (collected March–June 2015, $n = 103$), participants were quasi-randomized into treatment groups based on their availability to participate in the intervention immediately, and in the second wave (collected September 2015–February 2016, $n = 296$), participants were randomly allocated to treatment or wait-list. There were no significant differences in our independent variables (*MAOA* variant and trauma exposure; data on resilience were collected in the second wave only), enabling us to combine the two waves in order to increase sample size. In this paper, we focus on the 399 adolescents (221 males and 178 females) for whom biological samples were collected. The study received approval from the Prime Minister's Office of Jordan as well as ethical approval from Yale University (IRB ID 1502015359). Informed consent was obtained in Arabic from all participating adolescents and their parents. No participant who was asked to contribute a buccal sample refused to do so. See previous publications [57–59] for more details on the study method. All data analyzed in this study are presented in S1 and S2 Tables. All data are also available on Mendeley (<https://data.mendeley.com/datasets/>) doi:10.17632/xhs8tn7nkz.1).

Study population

Individuals with buccal samples and data on trauma exposure, resilience, and the outcome measures were chosen for study (T1 = 399) (Table 1). Some study participants were lost to follow-up at the T2 and T3 time points, resulting in a sample of 263 participants at T2 and 157 participants at T3. Because multilevel models are robust to partial data [60], individuals with missing time points were included in all analyses; thus, each data point contributes to the model.

Psychosocial measures

Lifetime trauma exposure was assessed with the Traumatic Events Checklist, adapted for use with adolescents in conflict regions [61]. Resilience was measured using the Child and Youth Resilience Measure (CYRM) that was specifically developed to assess resilience in youth living

Table 1. Sample characteristics of Syrian refugee participants.

	Male	Female
Sociodemographics		
# Participants at T1	221	178
# Participants at T2	136	127
# Participants at T3	78	79
Age, mean (SD) years	14.2 (1.81)	14.5 (1.88)
Genetic Data		
<i>MAOA-H</i> , # Homozygotes or Hemizygotes	125	50
<i>MAOA-H/L</i> , # Heterozygotes	NA	94
<i>MAOA-L</i> , # Homozygotes or Hemizygotes	96	34
Trauma exposure and Resilience Measures		
Lifetime Exposure to Traumatic Events* (SD)	6.76 (3.25)	5.81 (3.21)
Resilience, CYRM12 (SD)	49.8 (6.77)	49.1 (7.00)
Stress and Mental Health Measures		
Perceived Stress Scale, PSS* (SD)	28.1 (5.37)	29.4 (6.38)
Human Distress Scale, HD (SD)	39.7 (19.3)	44.5 (22.7)
Human Insecurity Scale, HI* (SD)	66.3 (20.0)	70.0 (19.2)
Arab Youth Mental Health Scale, AYMH* (SD)	34.6 (7.72)	37.2 (9.25)
Strengths & Difficulties Questionnaire, SDQ (SD)	14.9 (5.68)	16.3 (6.23)
Children’s Revised Impact of Events Scale, CRIES-8 (SD)	19.6 (10.6)	19.1 (11.7)

Values are reported as mean with standard deviation in parentheses and are from baseline. An asterisk (*) indicates significant differences between males and females at $p < 0.05$. For *MAOA*, repeat lengths 3.5, 4, and 5 were classified as high-activity variants and repeat lengths 2 and 3 were classified as low-activity variants. Since males have only one copy of the X-linked *MAOA* gene, there were no male heterozygotes. Resilience measures were collected for a subset of 163 males and 127 females.

<https://doi.org/10.1371/journal.pone.0219385.t001>

in adversity [55]. This measure of resilience is situated within a socioecological framework [55, 62] and assesses protective factors at individual, relational, and contextual levels, rather than treating resilience as an internal trait or using better-than-expected functioning as a proxy for resilience. Since the resilience measure was developed during this study, it was only collected in the second wave of survey with 163 males and 127 females.

We tested six psychosocial and mental health measures. The Perceived Stress Scale (PSS, 14 items) is a generalized measure of psychosocial stress used globally that has been validated in Jordan [63, 64]. Human Distress (HD, 12 items) is a measure of distress and Human Insecurity (HI, 10 items) is a measure of insecurity and captures feelings of fear in conflict-affected areas. Both HD and HI were developed for use in conflict settings in the Middle East and were validated in the West Bank [65]. The Arab Youth Mental Health scale (AYMH, 21 items) is used to screen for depression and anxiety in Arab youth [66]. The Strength and Difficulties Questionnaire (SDQ, 25 items) assesses behavioral and emotional mental health difficulties [67–69]. Finally, the Children's Revised Impact of Event Scale (CRIES-8, 8 items) measures symptoms of posttraumatic stress [70, 71].

Buccal sample collection and DNA extraction

Buccal samples were collected using Transport Swabs (APCO Laboratory Consumable Plastic, Jordan) or DNA Buccal Swabs (Isohelix, United Kingdom). Participants rinsed their mouths with water and then brushed both sides of their mouth with the collection swab for up to 30 seconds. DNA was extracted from the buccal swabs using either the Qiagen DNA Investigator Kit (Qiagen, USA) or Xtreme DNA Isolation Kit (Isohelix, United Kingdom). DNA extractions were performed according to manufacturer's recommendations with the exception that the AW2 wash was performed twice for swabs extracted using the Qiagen kit.

MAOA assay

Samples were PCR amplified based on a published protocol [34]. All samples used recommended reagent concentrations including 1 μ L of extracted DNA, Platinum Hot Start PCR Master Mix and Platinum GC Enhancer (Invitrogen, USA), and the following primer sequences: Forward 5' -CCC-AGG-CTG-CTC-CAG-AAA-C-3' and Reverse 5' -ACT-CAG-AAC-GGA-TGC-TCC-ATT-CG-3'. The PCR profile was as follows: 96°C for 10 min followed by 40 cycles of 94°C for 15 sec, 55°C for 15 sec, and 72°C for 30 sec and a final 3 min step at 72°C. Amplification reactions were electrophoresed on 2% agarose gels using Agarose SFR (VWR Life Science, USA) for 3 hours at 80V. Genotype completion rate was 97%. Genotyping discrepancy was determined by replicating approximately 1/3 of the samples ($n = 153$); only one sample showed a discrepancy and was typed a third time to resolve the discrepancy.

Statistical analyses

Each MAOA variant was classified according to transcriptional activity: repeat lengths of 3.5, 4, and 5 were classified as high-activity variants while repeat lengths of 2 and 3 were classified as low-activity variants. Since MAOA occurs on the X chromosome, males are hemizygous for MAOA, i.e. they carry only one MAOA variant. In statistical analyses, high-activity male hemizygotes and high-activity female homozygotes were given a score of 1, female heterozygotes with one high-activity and one low-activity variant were given a score of 0.5, and low-activity male hemizygotes and low-activity female homozygotes were given a score of 0. MAOA variant was treated as a categorical variable in males (0 or 1) and was treated as a continuous variable in females or analyses that included both males and females (0–1).

We used multilevel models (time points nested within participants) to test for effects of *MAOA* genotype, trauma exposure, and resilience on the psychosocial and mental health outcomes over the course of approximately one year. Multilevel models are robust to partially missing data, yielding higher statistical power than linear mixed models and regression analyses, while still taking advantage of a longitudinal design [60, 72]. These features are particularly useful for our data, given that more than half of the participants in this mobile refugee population were lost to follow-up at one year.

All statistical analyses were conducted using R [73]. Multilevel model analysis was conducted using linear mixed effects modeling via the nlme package with measurements at each time point (level one) nested within individuals (level two) [74]. Study site (Irbid, Jarash, Mafraq, Zarqa) and age at T1 were included as fixed effects, and these time-invariant covariates were included in all relevant analyses. Intervention program was treated as a time-varying covariate; all participants were given a score of 0 at T1, while at T2 and T3 those in the control group continued to receive a score of 0 (= has not received the intervention) and those in the treatment group received a score of 1 (= has received the intervention). Measurements of trauma exposure were treated as time-invariant and continuously distributed based on an individual's score at T1, consistent with previous studies using this sample [57, 75]. Measurements of resilience were treated as time-varying and measurements from all three time points were used. Models that included only Time were used to confirm that sufficient variation existed within the tested outcome to support multilevel modeling. Base models included time and covariates for age, collection site, and intervention. *MAOA*, trauma exposure, and resilience were added to the base models and were used to test for associations between the outcome measure and *MAOA*, trauma exposure, or resilience. Models for perceived psychosocial stress (PSS) were run in the following order (see Table 2):

Model 1. Testing for effects of *MAOA* variant on PSS:

- a. Time only
- b. Base (the base model includes time and the covariates: age, site, intervention)
- c. $PSS_{MAOA} = \text{Base} + MAOA + MAOA * \text{Time}$

Model 2. Testing for effects of *MAOA* variant and Trauma exposure on PSS:

- a. Time only
- b. Base (the base model includes time and the covariates: age, site, intervention)
- c. $PSS_{MAOA + Trauma} = \text{Base} + MAOA + MAOA * \text{Time} + \text{Trauma} + \text{Trauma} * \text{Time}$
- d. $PSS_{MAOA + Trauma + MAOA * Trauma} = \text{Base} + MAOA + MAOA * \text{Time} + \text{Trauma} + \text{Trauma} * \text{Time} + MAOA * \text{Trauma} + MAOA * \text{Trauma} * \text{Time}$

Model 3. Testing for effects of *MAOA* variant and Resilience on PSS:

- a. Time only
- b. Base (the base model includes time and the covariates: age, site, intervention)
- c. $PSS_{MAOA + Resilience} = \text{Base} + MAOA + MAOA * \text{Time} + \text{Resilience}$
- d. $PSS_{MAOA + Resilience + MAOA * Resilience} = \text{Base} + MAOA + MAOA * \text{Time} + \text{Resilience} + MAOA * \text{Resilience}$

Significance tests for all variables (as reported in Table 2) were conducted using ANOVAs that compared models with and without the variable being tested, i.e. in Model 1 outlined

Table 2. Perceived Stress Scale models with male participants.

Perceived Stress Scale (PSS)	Beta	Std. Error	p value
Model 1. Testing for effects of MAOA on PSS (n = 221)			
Time only	-0.074	0.016	5.6 x 10 ^{-6*}
PSS_{MAOA} = Base + MAOA + MAOA*Time			
MAOA	1.5	0.67	2.4 x 10 ⁻²
MAOA*Time	0.11	0.033	8.1 x 10 ^{-4*}
Model 2. Testing for effects of MAOA and Trauma exposure on PSS (n = 221)			
Time only	-0.074	0.016	5.6 x 10 ^{-6*}
PSS_{MAOA + Trauma} = Base + MAOA + MAOA*Time + Trauma + Trauma*Time			
MAOA	1.5	0.67	0.025
MAOA*Time	0.11	0.034	7.6 x 10 ^{-4*}
Trauma	0.14	0.11	0.19
Trauma*Time	0.00084	0.0058	0.88
PSS_{MAOA + Trauma + MAOA*Trauma} = Base + MAOA + MAOA*Time + Trauma + Trauma*Time + MAOA*Trauma + MAOA*Trauma*Time			
MAOA*Trauma	0.011	0.21	0.95
MAOA*Trauma*Time	0.0019	0.011	0.86
Model 3. Testing for effects of MAOA and Resilience on PSS (n = 163)			
Time only	-0.061	0.017	3.7 x 10 ^{-4*}
PSS_{MAOA} = Base + MAOA + MAOA*Time			
MAOA	2.2	0.78	6.1 x 10 ⁻³
MAOA*Time	0.11	0.035	1.4 x 10 ^{-3*}
PSS_{MAOA + Resilience} = Base + MAOA + MAOA*Time + Resilience			
MAOA	1.9	0.74	8.6 x 10 ⁻³
MAOA*Time	0.089	0.034	8.2 x 10 ⁻³
Resilience	-0.026	0.053	7.9 x 10 ^{-9*}
PSS_{MAOA + Resilience + MAOA*Resilience} = Base + MAOA + MAOA*Time + Resilience + MAOA*Resilience			
MAOA*Resilience	0.057	0.11	0.60

Time only models do not include covariates. Base models include time and covariates for age, study site, and intervention. Significance was tested individually for MAOA, Trauma exposure or Resilience with respect to association with the intercept of PSS (reported as MAOA or Trauma or Resilience) and for MAOA and Trauma exposure for association with the slope of PSS (reported as Trauma*Time or MAOA*Time). p values with an asterisk (*) remain significant after correction for multiple testing, i.e. $p < 1.67 \times 10^{-3}$.

<https://doi.org/10.1371/journal.pone.0219385.t002>

above, the significance of MAOA*Time was tested by comparing models with and without MAOA*Time. Significance after correction for multiple testing was calculated as $p < 0.05 / (5 \text{ psychosocial outcomes with sufficient variation for multilevel modeling} \times 2 \text{ sexes} \times 3 \text{ models}) = 1.67 \times 10^{-3}$.

In order to visualize the effects that we report, partial effect plots were constructed using the lme4, remef, and ggplot packages [76–78]. All plots were corrected for the same covariates (age, site, and intervention) that were included in the significance testing. Trauma exposure and resilience scores were treated as continuous variables in the multilevel models and when testing for significance, but were dichotomized when constructing the partial effect plots (Figs

1 and 2) to more clearly illustrate the effects (trauma exposure was dichotomized at <4 vs ≥ 4 events as this cutoff was previously found to be relevant for this sample [75] and resilience was dichotomized at the median based on averages of all resilience measures for each individual, i.e. median = 51).

Results

Sample characteristics

Sample characteristics for the Syrian refugee cohort are shown in Table 1. Genetic data were collected for 221 males and 178 females at T1, with retention of 66% and 39% of participants at T2 and T3, respectively. Exposure to traumatic events was higher for males relative to female participants, but levels of resilience were comparable by gender. No differences in trauma exposure or resilience levels were seen between *MAOA-L* and *MAOA-H* carriers. In terms of outcomes, psychosocial stress (PSS), insecurity (HI), and anxiety and depression (AYMH) symptoms were significantly higher in females.

Initial analyses by gender

We investigated six measures of psychosocial stress and mental health. All but one outcome (HI) had sufficient variation for multilevel modeling. With the cohort of males and females combined ($n = 399$), we found no significant associations between *MAOA* and the five tested outcomes (PSS, HD, AYMH, SDQ, or CRIES-8). We then divided the sample to examine males and females separately. In females ($n = 178$), no significant associations were found between *MAOA* and the five outcomes. In males ($n = 221$), associations were only significant for symptoms of psychosocial stress as measured on the Perceived Stress Scale (PSS); no associations were found between *MAOA* and either AYMH or SDQ, whilst there was insufficient variation for HD and CRIES-8 to test in multilevel models. Thus, all subsequent analyses focus on PSS in males.

MAOA

We found that *MAOA* was significantly associated with PSS over time, even after correction for multiple testing ($MAOA * Time p = 8.1 \times 10^{-4}$; Table 2, Model 1). Plots of *MAOA* and PSS show that *MAOA* variants exhibited different changes in PSS over time (Fig 1). Specifically, *MAOA-L* males had sharper reductions in PSS levels over time as compared to high *MAOA-H* males.

MAOA and trauma exposure

We next tested to see if the association identified above between *MAOA* and PSS was influenced by trauma exposure. Although previous studies have found interactive effects between *MAOA* and trauma, we found no interactive effect of *MAOA* and trauma exposure on PSS ($MAOA * Trauma p = 0.95$ and $MAOA * Trauma * Time p = 0.86$; Table 2, Model 2). As expected, *MAOA* was still directly associated with PSS over time, even with trauma exposure included in the model ($MAOA * Time p = 7.6 \times 10^{-4}$; Table 2, Model 2). Plots of *MAOA* and trauma exposure show that *MAOA-L* males had the sharpest reduction in PSS (Fig 2). Furthermore, there appeared to be an additive effect wherein *MAOA-L* males with low trauma exposure had the sharpest reductions in PSS over time, as compared to all *MAOA-H* males and *MAOA-L* males with high trauma exposure. In contrast to the difference in PSS shown by *MAOA-L* males with respect to trauma exposure, *MAOA-H* males with high and low trauma exposure showed the same PSS trajectory.




Fig 1. Partial effect plot of perceived psychosocial stress over time by MAOA variant in males. MAOA-L males had sharper reductions in levels of perceived stress (PSS) over time relative to MAOA-H males. Perceived stress symptom scores were adjusted for the effects of all covariates in the model and plotted (Y axis) for each participant at all time points (X axis).

<https://doi.org/10.1371/journal.pone.0219385.g001>

MAOA and resilience

We next tested to see if the association between MAOA and PSS was influenced by participants' levels of resilience. We first confirmed that the direct effect of MAOA on PSS over time reported above was also present in the smaller sample with resilience data ($MAOA \times Time$




Fig 2. Partial effect plot of perceived psychosocial stress over time by MAOA variant and trauma exposure in males. MAOA-L males had sharper reductions in levels of perceived stress (PSS) over time relative to MAOA-H males. Perceived stress symptom scores were adjusted for the effects of all covariates in the model and plotted (Y axis) for each participant at all time points (X axis). Partial effect plot lines were fitted for the four categories of MAOA variant (Low-activity/High-activity) and trauma exposure (Low/High). Trauma exposure was dichotomized around <4 vs ≥ 4 events in the figure for visualization purposes, but was treated as a continuous measure in multilevel models.

<https://doi.org/10.1371/journal.pone.0219385.g002>




Fig 3. Partial effect plot of perceived psychosocial stress over time by MAOA variant and resilience levels in males. MAOA-L males with high resilience had the sharpest reduction in levels of perceived stress (PSS) over time. Perceived stress symptom scores were adjusted for the effects of all covariates in the model and plotted (Y axis) for each participant at all time points (X axis). Partial effect plot lines were fitted for the four categories of MAOA variant (Low-activity/High-activity) and Resilience (Low/High). Resilience was dichotomized around the median (median = 51) in the figure purely for visualization purposes but was treated as a continuous measure in multilevel models.

<https://doi.org/10.1371/journal.pone.0219385.g003>

$p = 1.4 \times 10^{-3}$; Table 2, Model 3). When levels of resilience were added to the model, we found that MAOA was still associated with PSS, but no longer survived our conservative correction for multiple testing ($MAOA \times Time p = 8.2 \times 10^{-3}$; Table 2, Model 3). We also found that resilience was strongly associated with PSS (Resilience $p = 7.9 \times 10^{-9}$; Table 2, Model 3). We found no significant interactive effects between MAOA variants and resilience on PSS ($MAOA \times Resilience p = 0.28$; Table 2, Model 3). Plots show that higher levels of resilience were associated with lower PSS at baseline and with a sharper reduction in PSS over time (Fig 3). Furthermore, there appeared to be an additive effect with resilience wherein MAOA-L males with high resilience had the sharpest reductions in PSS over time compared to all MAOA-H males and MAOA-L males with low resilience (Fig 3).

Discussion

This is the first study to test the effect of early life trauma exposure, MAOA genetic variants, and a culturally-specific measure of resilience on psychosocial outcomes in a population of Syrian refugee youth over time. Specifically, we tested for direct and interactive effects of MAOA variants, trauma exposure, and resilience levels on psychosocial outcomes at baseline and over approximately one year. We tested a range of psychosocial and mental health outcomes in order to better understand the range of behaviors that may be influenced by trauma, MAOA, and resilience.

We found a significant association between MAOA and perceived psychosocial stress (PSS) over time, in males only, which remained significant after strict Bonferroni correction for multiple testing ($MAOA \times Time p = 8.1 \times 10^{-4}$; Table 2, Model 1). We found direct effects of MAOA on PSS, but no interactive effects between MAOA and trauma. We also found no associations between MAOA and the tested outcomes in the overall cohort, or in females only. These results suggest that MAOA plays a role in influencing levels of psychosocial stress in the males in our study population independent of their trauma exposure. Furthermore, MAOA was only

significantly associated with changes in PSS over time ($MAOA^*Time$ $p = 8.1 \times 10^{-4}$ vs $MAOA$ $p = 2.4 \times 10^{-2}$; Table 2). Specifically, *MAOA-L* males showed sharper reductions in PSS scores over time (Fig 1). The fact that the association of *MAOA* and PSS only emerges over time highlights the importance of conducting longitudinal research in order to understand the long-term effect of *MAOA* on psychosocial stress and possibly other measures of mental health.

We report no significant association between trauma and PSS in the males in our study population. However, we have found a significant association between trauma and PSS in the larger dataset of all Syrian refugees including those with no biological samples ($n = 446$) (unpublished data), so it may be that our smaller sample size of males only ($n = 221$) is not sufficient to detect an association between trauma and PSS. Interestingly, if we treat trauma as time-varying in the multilevel model, we do detect an association between trauma and PSS in males, although the p value ($p = 0.014$) is not significant after correction for multiple testing. This result suggests there is a trend of trauma exposure impacting PSS although the effect is not straightforward.

We also found that a culturally-relevant measure of resilience (CYRM-12) was strongly associated with psychosocial stress ($p = 7.9 \times 10^{-9}$; Table 2, Model 3). When resilience was added to the model, *MAOA* was no longer associated with PSS (after correction for multiple testing). Furthermore, we saw an additive effect with both trauma exposure and resilience levels wherein *MAOA-L* males with either low trauma exposure or high resilience showed the sharpest reductions in psychosocial stress (Figs 2 and 3).

Very few studies of complex phenotypes, such as response to trauma, include both genetic and psychosocial factors even though both genetic and environmental factors are involved in all complex phenotypes. By including both types of factors, we identified additive effects between *MAOA* and trauma exposure and between *MAOA* and resilience on psychosocial stress. Additive effects act independently, and not as gene x environment interaction effects wherein the environmental factor exerts different effects on an individual based on his or her genotype. Furthermore, by comparing the effects of *MAOA* and resilience, we conclude that *MAOA* variation impacts levels of psychosocial stress in males, but that resilience appears to be a more influential factor.

Our study is the second to examine the effect of *MAOA* and resilience on mental health outcomes. We find an additive effect wherein *MAOA-L* males with high resilience showed the sharpest reductions in perceived psychosocial stress (Fig 3). The only other study to investigate the impact of protective factors and *MAOA* variants on mental health found a similar result with *MAOA-L* carriers. Specifically, Kinnally et al. [40] found that high perceived parental care mitigated the effect of a childhood stressor on impulsivity scores in *MAOA-L* females (males were not studied); level of perceived parental care had no effect on impulsivity in homozygous *MAOA-H* females.

Our results are consistent with Pluess and Belsky's model of environmental sensitivity that posits carriers of sensitivity variants are more sensitive to both positive and negative environmental exposures [52, 53, 79]. Pluess [51] proposed a theoretical framework in which environmental sensitivity is a function of specific genetic variants, and genetic factors predict individual differences in environmental sensitivity to both positive and adverse conditions. Belsky and Pluess and colleagues [80, 81] specifically proposed that due to genetic differences, some children would benefit more from intervention efforts and that evaluations of intervention efficacy should account for such genetic variation. In the case of *MAOA*, our results suggest that *MAOA-L* variants may function as sensitivity variants and carriers may be more responsive to the effect of both negative (trauma exposure) and positive (levels of resilience) factors. Similarly, other studies have found that the *MAOA-L* variant is most sensitive to a positive factor (high perceived parental care; [40]) and to negative factors (aggressive and

antisocial behavior in the presence of childhood trauma [17, 25, 82, 83]). Furthermore, in a brain imaging study, Buckholtz et al. [28] proposed that *MAOA-L* individuals may be more susceptible to environmental factors than *MAOA-H* individuals due to their amygdala dysregulation and compensatory response between the amygdala and the ventromedial prefrontal cortex. It is important to note that Pluess [51] posited that sensitivity variants act through a gene x environment mechanism with no proposed difference between genetic variants in the absence of adverse or positive conditions. We found a direct effect of *MAOA* on PSS, rather than gene x environment effects with trauma exposure or resilience, which may suggest that sensitivity variants can also act through an additive mechanism.

As noted above, we only identified associations of *MAOA* and resilience with PSS in males. This result is consistent with other studies that have found the effect of *MAOA* on measures of psychosocial health and behavior to be weak or non-existent in females [28, 39, 83, 84]. Since *MAOA* occurs on the X chromosome, males and females differ in copy number of *MAOA* and random X chromosome inactivation in females means that females likely have variable *MAOA* levels throughout the body relative to males. Studies of brain morphology and brain activity have found different effects of childhood trauma on male and female brains [23, 27] so *MAOA* may have different biological effects in males and females.

There are three main limitations to note in our study. First, we focused on a single gene. *MAOA* is a well-supported candidate gene to investigate the impact of childhood trauma, but additional genes should be studied in order to fully understand the impact of childhood trauma and resilience on adult behaviors, particularly in refugee populations with exposure to high levels of trauma. Second, our focus on Syrian refugees, who have relatively high levels of trauma exposure, may have restricted the level of population variation in some of the psychosocial health outcomes. Third, our sample size is small, particularly since the association we identified was only found in males, which reduced our sample size. A recent meta-analysis found no support for 18 previously identified candidate genes for depression, suggesting that previous studies had been underpowered [85]. Research on high trauma populations can help us understand the impact of trauma on mental health, but collection of a large number of longitudinal samples from a highly mobile refugee population will always be challenging [86]. The fact that, even after correction for multiple testing, we detected significant associations with psychosocial stress in our study population is encouraging. Nevertheless, work with a larger, more heterogeneous sample is needed to more fully evaluate the effects of *MAOA*, trauma exposure, and resilience on psychosocial stress and mental health measures. In order to fully understand the impact of trauma on mental health, more work with vulnerable populations is necessary, particularly in non-Western contexts with high levels of trauma exposure.

Conclusions

We report significant effects of *MAOA* genetic variants and resilience levels on changes in perceived psychosocial stress over time, in a population of Syrian refugee male youth. We also find an additive effect wherein males with low-activity *MAOA* variants who reported low trauma exposure or high resilience had the sharpest reductions in perceived psychosocial stress over time. Our results highlight the potential scientific value of careful and ethical collection of data and biological samples in populations with high levels of trauma exposure. In contexts of war, violence, and forced migration, addressing the root causes of human suffering and the health consequences of toxic stress and trauma exposure is of vital importance. It is also prudent to reach for an in-depth understanding of the interplay between genetic and psychosocial factors that are able to influence the expression of interpersonal and social behaviors over the life course.

Supporting information

S1 Table. Data on trauma exposure, resilience level, psychosocial stress and mental health measures, and MAOA variants for all study participants. See [S2 Table](#) for codebook for variable names presented here.

(CSV)

S2 Table. Codebook for variable names presented in [S1 Table](#).

(XLSX)

Acknowledgments

We gratefully acknowledge the Syrian families who participated in this project. We thank Jane Macphail, Director of the Advancing Adolescents programme implemented in Jordan as part of the No Lost Generation initiative in the wider region; Noura Shaed, Natasha Shawarib and Jon Kurtz at Mercy Corps; and the field team affiliated to the organization Taghyeer, especially Dima Hamadmad, Ghufraan Abydayyeh, Rahmeh Hyari and Sana'a Bakeer.

Author Contributions

Conceptualization: Rana Dajani, Catherine Panter-Brick, Connie J. Mulligan.

Data curation: Catherine Panter-Brick.

Formal analysis: Christopher J. Clukay, Kristin Hadfield.

Investigation: Christopher J. Clukay, Kristin Hadfield.

Methodology: Christopher J. Clukay, Jacklyn Quinlan.

Project administration: Rana Dajani, Catherine Panter-Brick.

Resources: Rana Dajani.

Supervision: Rana Dajani, Catherine Panter-Brick, Connie J. Mulligan.

Visualization: Christopher J. Clukay.

Writing – original draft: Christopher J. Clukay, Kristin Hadfield, Connie J. Mulligan.

Writing – review & editing: Christopher J. Clukay, Rana Dajani, Kristin Hadfield, Catherine Panter-Brick, Connie J. Mulligan.

References

1. Edwards VJ, Holden GW, Felitti VJ, Anda RF. Relationship between multiple forms of childhood maltreatment and adult mental health in community respondents: Results from the adverse childhood experiences study. *American Journal of Psychiatry*. 2018; 160(8):1453–60.
2. Keyes KM, Eaton NR, Krueger RF, McLaughlin KA, Wall MM, Grant BF, et al. Childhood maltreatment and the structure of common psychiatric disorders. *British Journal of Psychiatry*. 2012; 200(2):107–15. Epub 2018/01/02. <https://doi.org/10.1192/bjp.bp.111.093062> PMID: 22157798
3. Scott KM, McLaughlin KA, Smith DAR, Ellis PM. Childhood maltreatment and DSM-IV adult mental disorders: Comparison of prospective and retrospective findings. *British Journal of Psychiatry*. 2012; 200(6):469–75. <https://doi.org/10.1192/bjp.bp.111.103267> PMID: 22661679
4. Clemens V, Huber-Lang M, Plener PL, Brähler E, Brown RC, Fegert JM. Association of child maltreatment subtypes and long-term physical health in a German representative sample. *European Journal of Psychotraumatology*. 2018; 9(1):1510278. <https://doi.org/10.1080/20008198.2018.1510278> PMC6136347. PMID: 30220980

5. Thompson MP, Kingree JB, Lamis D. Associations of adverse childhood experiences and suicidal behaviors in adulthood in a U.S. nationally representative sample. *Child: Care, Health and Development*. 2018;2018(0):1–8. <https://doi.org/10.1111/cch.12617> PMID: 30175459
6. Stepanikova I, Baker E, Oates G, Acharya S, Uddin J, Thon V, et al. Perinatal Maternal Stress and Susceptibility to Infectious Diseases in Later Childhood: An Early Life Programming Perspective. *J Psychol*. 2018;in press. <https://doi.org/10.1080/00223980.2018.1483311> PMID: 30265824
7. McDade TW, Ryan C, Jones MJ, Maclsaac JL, Morin AM, Meyer JM, et al. Social and physical environments early in development predict DNA methylation of inflammatory genes in young adulthood. *Proceedings of the National Academy of Sciences*. 2017; 114(29):7611–6. <https://doi.org/10.1073/pnas.1620661114> PMID: 28673994
8. Obradović J, Boyce WT. Individual Differences in Behavioral, Physiological, and Genetic Sensitivities to Contexts: Implications for Development and Adaptation. *Developmental Neuroscience*. 2009; 31(4):300–8. <https://doi.org/10.1159/000216541> PMID: 19546567
9. SaveTheChildren. Invisible wounds: The impact of six years of war on the mental health of Syria's children. 2018. Available from: <https://www.savethechildren.org/content/dam/global/reports/emergency-humanitarian-response/invisible-wounds.pdf>. Accessed 13.07.2018.
10. UNHCR. Syrian regional refugee response in Jordan 2017. Available from: <https://data.unhcr.org/syrianrefugees/country.php?id=107>.
11. Masten AS. Global Perspectives on Resilience in Children and Youth. *Child Development*. 2014; 85(1):6–20. <https://doi.org/10.1111/cdev.12205> PMID: 24341286
12. Reed RV, Fazel M, Jones L, Panter-Brick C, Stein A. Mental health of displaced and refugee children resettled in low-income and middle-income countries: risk and protective factors. *The Lancet*. 2012; 379(9812):250–65. [https://doi.org/10.1016/S0140-6736\(11\)60050-0](https://doi.org/10.1016/S0140-6736(11)60050-0).
13. UNHCR. Global trends: Forced displacement in 2015: Geneva, Switzerland: UNHCR; 2016.
14. UNHCR. Syria Regional Refugee Response 2018. Available from: <https://www.savethechildren.org/content/dam/global/reports/emergency-humanitarian-response/invisible-wounds.pdf>. Accessed 13.07.2018.
15. Sabol SZ, Hu S, Hamer D. A functional polymorphism in the monoamine oxidase A gene promoter. *Human Genetics*. 1998; 103:273–9. PMID: 9799080
16. Brunner H, Nelen M, Breakefield X, Ropers H, van Oost B. Abnormal behavior associated with a point mutation in the structural gene for monoamine oxidase A. *Science*. 1993; 262(5133):578–80. <https://doi.org/10.1126/science.8211186> PMID: 8211186
17. Caspi A, McClay J, Moffitt TE, Mill J, Martin J, Craig IW, et al. Role of Genotype in the Cycle of Violence in Maltreated Children. *Science*. 2002; 297(5582):851–4. <https://doi.org/10.1126/science.1072290> PMID: 12161658
18. Lee SS. Deviant Peer Affiliation and Antisocial Behavior: Interaction with Monoamine Oxidase A (MAOA) Genotype. *Journal of Abnormal Child Psychology*. 2011; 39(3):321–32. <https://doi.org/10.1007/s10802-010-9474-2> PMID: 21152968
19. Mertins V, Schote AB, Hoffeld W, Griessmair M, Meyer J. Genetic Susceptibility for Individual Cooperation Preferences: The Role of Monoamine Oxidase A Gene (MAOA) in the Voluntary Provision of Public Goods. *PLOS ONE*. 2011; 6(6):e20959. <https://doi.org/10.1371/journal.pone.0020959> PMID: 21698196
20. Frazzetto G, Di Lorenzo G, Carola V, Proietti L, Sokolowska E, Siracusano A, et al. Early Trauma and Increased Risk for Physical Aggression during Adulthood: The Moderating Role of MAOA Genotype. *PLoS ONE*. 2007; 2(5):e486. <https://doi.org/10.1371/journal.pone.0000486> PMID: 17534436
21. Stetler DA, Davis C, Leavitt K, Schriger I, Benson K, Bhakta S, et al. Association of low-activity MAOA allelic variants with violent crime in incarcerated offenders. *Journal of Psychiatric Research*. 2014; 58:69–75. <https://doi.org/10.1016/j.jpsychires.2014.07.006> PMID: 25082653
22. Tikkanen R, Ducci F, Goldman D, Holli M, Lindberg N, Tiihonen J, et al. MAOA Alters the Effects of Heavy Drinking and Childhood Physical Abuse on Risk for Severe Impulsive Acts of Violence Among Alcoholic Violent Offenders. *Alcoholism: Clinical and Experimental Research*. 2010; 34(5):853–60. <https://doi.org/10.1111/j.1530-0277.2010.01157.x> PMID: 20201935
23. Kolla NJ, Patel R, Meyer JH, Chakravarty MM. Association of monoamine oxidase-A genetic variants and amygdala morphology in violent offenders with antisocial personality disorder and high psychopathic traits. *Scientific Reports*. 2017; 7(1):9607. <https://doi.org/10.1038/s41598-017-08351-w> PMID: 28851912
24. Tielbeek JJ, Medland SE, Benyamin B, Byrne EM, Heath AC, Madden PAF, et al. Unraveling the Genetic Etiology of Adult Antisocial Behavior: A Genome-Wide Association Study. *PLOS ONE*. 2012; 7(10):e45086. <https://doi.org/10.1371/journal.pone.0045086> PMID: 23077488

25. Byrd AL, Manuck SB. MAOA, Childhood Maltreatment, and Antisocial Behavior: Meta-analysis of a Gene-Environment Interaction. *Biological Psychiatry*. 2014; 75(1):9–17. <https://doi.org/10.1016/j.biopsych.2013.05.004> PMID: 23786983
26. Schmidt LG, Sander T, Kuhn S, Smolka M, Rommelspacher H, Samochowicz J, et al. Different allele distribution of a regulatory MAOA gene promoter polymorphism in antisocial and anxious-depressive alcoholics. *Journal of Neural Transmission*. 2000; 107(6):681–9. <https://doi.org/10.1007/s007020070069> PMID: 10943908
27. Holz N, Boecker R, Buchmann AF, Blomeyer D, Baumeister S, Hohmann S, et al. Evidence for a Sex-Dependent MAOA \times Childhood Stress Interaction in the Neural Circuitry of Aggression. *Cerebral Cortex*. 2016; 26(3):904–14. <https://doi.org/10.1093/cercor/bhu249> PMID: 25331606
28. Buckholtz JW, Callicott JH, Kolachana B, Hariri AR, Goldberg TE, Genderson M, et al. Genetic variation in MAOA modulates ventromedial prefrontal circuitry mediating individual differences in human personality. *Molecular Psychiatry*. 2008; 13:313–24. <https://doi.org/10.1038/sj.mp.4002020> <https://www.nature.com/articles/4002020#supplementary-information>. PMID: 17519928
29. Sjöberg RL, Ducci F, Barr CS, Newman TK, Dell'Osso L, Virkkunen M, et al. A Non-Additive Interaction of a Functional MAO-A VNTR and Testosterone Predicts Antisocial Behavior. *Neuropsychopharmacology*. 2007; 33:425. <https://doi.org/10.1038/sj.npp.1301417> PMID: 17429405
30. Andersen SL, Tomada A, Vincow ES, Valente E, Polcari A, Teicher MH. Preliminary Evidence for Sensitive Periods in the Effect of Childhood Sexual Abuse on Regional Brain Development. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 2008; 20(3):292–301. <https://doi.org/10.1176/appi.neuropsych.20.3.292> PMID: 18806232.
31. Fox SE, Levitt P, Nelson III CA. How the Timing and Quality of Early Experiences Influence the Development of Brain Architecture. *Child Development*. 2010; 81(1):28–40. <https://doi.org/10.1111/j.1467-8624.2009.01380.x> PMID: 20331653
32. Giedd JN. Structural Magnetic Resonance Imaging of the Adolescent Brain. *Annals of the New York Academy of Sciences*. 2004; 1021(1):77–85. <https://doi.org/10.1196/annals.1308.009> PMID: 15251877
33. Tottenham N, Sheridan M. A review of adversity, the amygdala and the hippocampus: a consideration of developmental timing. *Frontiers in Human Neuroscience*. 2010; 3(68). <https://doi.org/10.3389/neuro.09.068.2009> PMID: 20161700
34. Enoch M-A, Steer CD, Newman TK, Gibson N, Goldman D. Early life stress, MAOA, and gene-environment interactions predict behavioral disinhibition in children. *Genes, Brain and Behavior*. 2010; 9(1):65–74. <https://doi.org/10.1111/j.1601-183X.2009.00535.x> PMID: 19804559
35. McGregor NW, Hemmings SMJ, Erdman L, Calmarza-Font I, Stein DJ, Lochner C. Modification of the association between early adversity and obsessive-compulsive disorder by polymorphisms in the MAOA, MAOB and COMT genes. *Psychiatry Research*. 2016; 246:527–32. <https://doi.org/10.1016/j.psychres.2016.10.044> PMID: 27821364
36. Caporaso N, Gu F, Chatterjee N, Sheng-Chih J, Yu K, Yeager M, et al. Genome-Wide and Candidate Gene Association Study of Cigarette Smoking Behaviors. *PLOS ONE*. 2009; 4(2):e4653. <https://doi.org/10.1371/journal.pone.0004653> PMID: 19247474
37. Zhong S, Israel S, Xue H, Ebbstein RP, Chew SH. Monoamine Oxidase A Gene (MAOA) Associated with Attitude Towards Longshot Risks. *PLOS ONE*. 2010; 4(12):e8516. <https://doi.org/10.1371/journal.pone.0008516> PMID: 20046877
38. Lundwall RA, Watkins JK. Genetic Influence on Slope Variability in a Childhood Reflexive Attention Task. *PLOS ONE*. 2015; 10(6):e0130668. <https://doi.org/10.1371/journal.pone.0130668> PMID: 26102342
39. Goldman D, Rosser AA. MAOA–Environment Interactions: Results May Vary. *Biological Psychiatry*. 2014; 75(1):2–3. <https://doi.org/10.1016/j.biopsych.2013.09.015> PMID: 24314060
40. Kinnally EL, Huang Y-y, Haverly R, Burke AK, Galfalvy H, Brent DP, et al. Parental care moderates the influence of MAOA-uVNTR genotype and childhood stressors on trait impulsivity and aggression in adult women. *Psychiatric Genetics*. 2009; 19(3):126–33. <https://doi.org/10.1097/YPG.0b013e32832a50a7> 00041444-200906000-00003. PMID: 19357553
41. Aro H. Risk and protective factors in depression: a developmental perspective. *Acta Psychiatrica Scandinavica*. 1994; 89(s377):59–64. <https://doi.org/10.1111/j.1600-0447.1994.tb05804.x>
42. Betancourt TS, Khan KT. The mental health of children affected by armed conflict: Protective processes and pathways to resilience. *International Review of Psychiatry*. 2008; 20(3):317–28. <https://doi.org/10.1080/09540260802090363> PMID: 18569183
43. Doom JR, Gunnar MR. Stress physiology and developmental psychopathology: Past, present, and future. *Development and Psychopathology*. 2013; 25(4pt2):1359–73. Epub 2013/12/17. <https://doi.org/10.1017/S0954579413000667> PMID: 24342845

44. Masten AS, Narayan AJ. Child Development in the Context of Disaster, War, and Terrorism: Pathways of Risk and Resilience. *Annual Review of Psychology*. 2012; 63(1):227–57. <https://doi.org/10.1146/annurev-psych-120710-100356> PMID: 21943168.
45. Masten A, Barnes A. Resilience in Children: Developmental Perspectives. *Children*. 2018; 5(7):98. <https://doi.org/10.3390/children5070098> PMID: 30018217
46. Guan S-SA, Bower JE, Almeida DM, Cole SW, Dahl RE, Irwin MR, et al. Parental support buffers the association of depressive symptoms with cortisol and C-reactive protein during adolescence. *Brain, Behavior, and Immunity*. 2016; 57:134–43. <https://doi.org/10.1016/j.bbi.2016.03.007> PMID: 26995316
47. Labella MH, Kalstabbakken A, Johnson J, Leppa J, Robinson N, Masten AS, et al. Promoting Resilience by Improving Children's Sleep: Feasibility Among Families Living in Supportive Housing. *Progress in Community Health Partnerships: Research, Education, and Action*. 2017; 11(3):285–93. <https://doi.org/10.1353/cpr.2017.0033> PMID: 29056620
48. Koni E, Moradi S, Arahanga-Doyle H, Neha T, Hayhurst JG, Boyes M, et al. Promoting resilience in adolescents: A new social identity benefits those who need it most. *PLOS ONE*. 2019; 14(1):e0210521. <https://doi.org/10.1371/journal.pone.0210521> PMID: 30629716
49. Belsky J, Conger R, Capaldi DM. The intergenerational transmission of parenting: Introduction to the special section. *Developmental Psychology*. 2009; 45(5):1201–4. <https://doi.org/10.1037/a0016245> PMID: 19702385
50. Shaffer A, Burt KB, Obradović J, Herbers JE, Masten AS. Intergenerational continuity in parenting quality: The mediating role of social competence. *Developmental Psychology*. 2009; 45(5):1227–40. <https://doi.org/10.1037/a0015361> PMID: 19702388
51. Pluess M. Vantage Sensitivity: Environmental Sensitivity to Positive Experiences as a Function of Genetic Differences. *Journal of Personality*. 2017; 85(1):38–50. <https://doi.org/10.1111/jopy.12218> PMID: 26271007
52. Pluess M, Belsky J. Vantage sensitivity: Individual differences in response to positive experiences. *Psychological Bulletin*. 2013; 139(4):901–16. <https://doi.org/10.1037/a0030196> PMID: 23025924
53. Hartman S, Belsky J. Prenatal stress and enhanced developmental plasticity. *Journal of Neural Transmission*. 2018. <https://doi.org/10.1007/s00702-018-1926-9> PMID: 30206701
54. Buckholtz JW, Callicott JH, Kolachana B, Hariri AR, Goldberg TE, Genderson M, et al. Genetic variation in MAOA modulates ventromedial prefrontal circuitry mediating individual differences in human personality. *Molecular Psychiatry*. 2007; 13:313. <https://doi.org/10.1038/sj.mp.4002020> <https://www.nature.com/articles/4002020#supplementary-information>. PMID: 17519928
55. Panter-Brick C, Hadfield K, Dajani R, Eggerman M, Ager A, Ungar M. Resilience in Context: A Brief and Culturally Grounded Measure for Syrian Refugee and Jordanian Host-Community Adolescents. *Child Development*. 2017:Epub ahead of print. <https://doi.org/10.1111/cdev.12868> PMID: 28617937
56. Meyer-Lindenberg A, Buckholtz JW, Kolachana B, R. Hariri A, Pezawas L, Blasi G, et al. Neural mechanisms of genetic risk for impulsivity and violence in humans. *Proceedings of the National Academy of Sciences*. 2006; 103(16):6269–74. <https://doi.org/10.1073/pnas.0511311103> PMID: 16569698
57. Dajani R, Hadfield K, van Uum S, Greff M, Panter-Brick C. Hair cortisol concentrations in war-affected adolescents: A prospective intervention trial. *Psychoneuroendocrinology*. 2018; 89:138–46. <https://doi.org/10.1016/j.psyneuen.2017.12.012> PMID: 29358120
58. Panter-Brick C, Kurtz J, Dajani R. What strong partnerships achieve: innovations in research and practice. *Humanitarian Exchange*. 2018; 72:15–9.
59. Panter-Brick C, Wiley K, Sancilio A, Dajani R, Hadfield K. C-reactive protein, Epstein-Barr virus, and cortisol trajectories in refugee and non-refugee youth: Links with stress, mental health, and cognitive function during a randomized controlled trial. *Brain, Behavior, and Immunity*. 2019. <https://doi.org/10.1016/j.bbi.2019.02.015>.
60. Curran PJ, Obeidat K, Losardo D. Twelve Frequently Asked Questions About Growth Curve Modeling. *Journal of cognition and development: official journal of the Cognitive Development Society*. 2010; 11(2):121–36. <https://doi.org/10.1080/15248371003699969> PMC3131138. PMID: 21743795
61. Panter-Brick C, Eggerman M, Gonzalez V, Safdar S. Violence, suffering, and mental health in Afghanistan: a school-based survey. *Lancet*. 2009; 374(9692):807–16. Epub 2009/08/25. [https://doi.org/10.1016/S0140-6736\(09\)61080-1](https://doi.org/10.1016/S0140-6736(09)61080-1) PMID: 19699514; PubMed Central PMCID: PMC2748901.
62. Ungar M, Liebenberg L. Assessing Resilience Across Cultures Using Mixed Methods: Construction of the Child and Youth Resilience Measure. *Journal of Mixed Methods Research*. 2011; 5(2):126–49. <https://doi.org/10.1177/1558689811400607>
63. Almadi T, Cathers I, Hamdan Mansour AM, Chow CM. An Arabic version of the Perceived Stress Scale: Translation and validation study. *International Journal of Nursing Studies*. 2012; 49(1):84–9. <https://doi.org/10.1016/j.ijnurstu.2011.07.012> PMID: 21851941

64. Cohen S, Kamarck T, Mermelstein R. A Global Measure of Perceived Stress. *Journal of Health and Social Behavior*. 1983; 24(4):385–96. <https://doi.org/10.2307/2136404> PMID: 6668417
65. Ziadni M, Hammoudeh W, Rmeileh NMEA, Hogan D, Shannon H, Giacaman R. Sources of Human Insecurity in Post-War Situations: The Case of Gaza. *Journal of human security*. 2011; 7(3):10.3316/JHS0703023. <https://doi.org/10.3316/JHS0703023> PMID: 24416084.
66. Mahfoud Z, Abdulrahim S, Taha MB, Harpham T, El Hajj T, Makhoul J, et al. Validation of the Arab Youth Mental Health scale as a screening tool for depression/anxiety in Lebanese children. *Child and Adolescent Psychiatry and Mental Health*. 2011; 5:9–. <https://doi.org/10.1186/1753-2000-5-9> PMC3070665. PMID: 21435213
67. Almaqrami MH, Shuwail AY. Validity of the self-report version of the strengths and difficulties questionnaire in Yemen. *Saudi Medical Journal*. 2004; 25(5):592–601. PMID: 15138526
68. Alyahri A, Goodman R. Validation of the Arabic Strengths and Difficulties Questionnaire and the Development and Well-Being Assessment. *Eastern Mediterranean Health Journal*. 2006; 12 Suppl 2:S138–S46.
69. Goodman A, Goodman R. Strengths and Difficulties Questionnaire as a Dimensional Measure of Child Mental Health. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2009; 48(4):400–3. <https://doi.org/10.1097/CHI.0b013e3181985068>.
70. Punamäki R-L, Palosaari E, Diab M, Peltonen K, Qouta SR. Trajectories of posttraumatic stress symptoms (PTSS) after major war among Palestinian children: Trauma, family- and child-related predictors. *Journal of Affective Disorders*. 2015; 172:133–40. <https://doi.org/10.1016/j.jad.2014.09.021> PMID: 25451407
71. Veronese G, Pepe A. Psychometric properties of IES-R, short Arabic version in contexts of military violence. *Research on Social Work Practice*. 2013; 23(710–718).
72. Muthén BO, Curran PJ. General longitudinal modeling of individual differences in experimental designs: A latent variable framework for analysis and power estimation. *Psychological Methods*. 1997; 2(4):371–402. <https://doi.org/10.1037/1082-989X.2.4.371>
73. Team RC. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2015.
74. Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC. nlme: Linear and nonlinear mixed effects models. 3.1–137 ed; 2016.
75. Panter-Brick C, Dajani R, Eggerman M, Hermsilla S, Sancilio A, Ager A. Insecurity, distress and mental health: experimental and randomized controlled trials of a psychosocial intervention for youth affected by the Syrian crisis. *J Child Psychol Psychiatry*. 2018; 59(5):523–41. Epub 2017/10/03. <https://doi.org/10.1111/jcpp.12832> PMID: 28967980.
76. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*. 2015; 67(1):1–48. <https://doi.org/10.18637/jss.v067.i01>
77. Hohenstein S, Kliegl R. remef: Remove partial effects. 1.0.6.9000 ed; 2018.
78. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag; 2009.
79. Pluess M, Assary E, Lionetti F, Lester KJ, Krapohl E, Aron EN, et al. Environmental sensitivity in children: Development of the Highly Sensitive Child Scale and identification of sensitivity groups. *Developmental Psychology*. 2018; 54(1):51–70. <https://doi.org/10.1037/dev0000406> PMID: 28933890
80. Belsky J, van Ijzendoorn MH. Genetic differential susceptibility to the effects of parenting. *Current Opinion in Psychology*. 2017; 15:125–30. <https://doi.org/10.1016/j.copsyc.2017.02.021> PMID: 28813251
81. Pluess M, Boniwell I. Sensory-Processing Sensitivity predicts treatment response to a school-based depression prevention program: Evidence of Vantage Sensitivity. *Personality and Individual Differences*. 2015; 82:40–5. <https://doi.org/10.1016/j.paid.2015.03.011>.
82. Ficks CA, Waldman ID. Candidate Genes for Aggression and Antisocial Behavior: A Meta-analysis of Association Studies of the 5HTTLPR and MAOA-uVNTR. *Behavior Genetics*. 2014; 44(5):427–44. <https://doi.org/10.1007/s10519-014-9661-y> PMID: 24902785
83. Kim-Cohen J, Caspi A, Taylor A, Williams B, Newcombe R, Craig IW, et al. MAOA, maltreatment, and gene–environment interaction predicting children’s mental health: new evidence and a meta-analysis. *Molecular Psychiatry*. 2006; 11:903. <https://doi.org/10.1038/sj.mp.4001851> PMID: 16801953
84. Guo G, Ou X-M, Roettger M, Shih JC. The VNTR 2 repeat in MAOA and delinquent behavior in adolescence and young adulthood: associations and MAOA promoter activity. *European Journal Of Human Genetics*. 2008; 16:626. <https://doi.org/10.1038/sj.ejhg.5201999> PMID: 18212819
85. Border R, Johnson EC, Evans LM, Smolen A, Berley N, Sullivan PF, et al. No Support for Historical Candidate Gene or Candidate Gene-by-Interaction Hypotheses for Major Depression Across Multiple Large Samples. *American Journal of Psychiatry*. 2019; 176(5):376–87. <https://doi.org/10.1176/appi.ajp.2018.18070881> PMID: 30845820.

86. Moghames P, McEwen F, Pluess M. Mental health research among Syrian refugees in Lebanon: challenges and solutions. *Humanitarian Exchange* [Internet]. 2018; 72:[39–42 pp.]. Available from: <https://odihpn.org/magazine/mental-health-research-among-syrian-refugees-in-lebanon-challenges-and-solutions/>.